

3) 巡回調査成績

(1) 稲

ア 早期水稻

調査時期			葉いもち				紋枯病				ばか苗病				萎縮病			
年	月	期	調査発生		発病株率		調査発生		発病株率		調査発生		発病株率		調査発生		発病株率	
			調査数	発生数	最高	平均	調査数	発生数	最高	平均	調査数	発生数	最高	平均	調査数	発生数	最高	平均
98	5	前	20	0	0	0	20	0	0	0	20	0	0	0	20	0	0	0
		後	42	0	0	0	42	0	0	0	42	0	0	0	42	0	0	0
	6	前	15	0	0	0	15	0	0	0	15	0	0	0	15	0	0	0
		後	38	5.3	1	0.01	38	0	0	0	38	0	0	0	38	0	0	0
	7	前	13	7.7	10	0.8	13	15.4	2	0.3	13	0	0	0	13	0	0	0
		後	33	21.2	16	1.8	33	21.2	12	1.9	33	3	1	0.03	33	0	0	0
	8	前	-	-	-	-	12	75	70	12.8	-	-	-	-	12	0	0	0
		後	-	-	-	-	17	58.8	92	18.2	-	-	-	-	17	0	0	0

調査時期			稲葉枯病				白葉枯病				こま葉枯病				穂いもち							
年	月	期	調査発生		発病株率		調査発生		発病株率		調査発生		発病株率		調査発生		発病株率		発病率		発病率	
			調査数	発生数	最高	平均	調査数	発生数	最高	平均	調査数	発生数	最高	平均	調査数	発生数	最高	平均	最高	平均	最高	平均
98	5	前	20	0	0	0	20	0	0	0	-	-	-	-	-	-	-	-	-	-	-	-
		後	42	0	0	0	42	0	0	0	-	-	-	-	-	-	-	-	-	-	-	-
	6	前	15	0	0	0	15	0	0	0	15	0	0	0	-	-	-	-	-	-	-	-
		後	38	0	0	0	38	0	0	0	38	0	0	0	-	-	-	-	-	-	-	-
	7	前	13	7.7	4	0.3	13	0	0	0	13	0	0	0	13	0	0	0	0	0	0	0
		後	33	0	0	0	33	0	0	0	33	12.1	100	5.2	33	0	0	0	0	0	0	0
	8	前	12	0	0	0	12	0	0	0	12	8.3	6	0.5	12	0	0	0	0	0	0	0
		後	17	0	0	0	17	0	0	0	17	0	0	0	17	0	0	0	0	0	0	0

調査時期			もみ枯細菌病						ツマグロヨコバイ									
年	月	期	調査発生		発病株率		発病率		調査発生		寄生株率		株当成虫数		株当幼虫数		合計虫数	
			調査数	発生数	最高	平均	最高	平均	調査数	発生数	最高	平均	最高	平均	最高	平均	最高	平均
98	5	前	-	-	-	-	-	-	20	0	0	0	0	0	0	0	0	0
		後	-	-	-	-	-	-	42	7	12	1	0.8	0.03	0.1	0.003	0.9	0.03
	6	前	-	-	-	-	-	-	15	80	32	9.9	0.08	0.01	0.4	0.1	0.4	0.1
		後	-	-	-	-	-	-	38	21.2	52	5.6	0.3	0.01	1	0.1	1	0.1
	7	前	13	0	0	0	0	0	13	61.5	60	13.8	0.5	0.05	1.5	0.3	1.5	0.3
		後	33	0	0	0	0	0	33	36.4	72	6.7	0.7	0.03	1.2	0.1	1.9	0.1
	8	前	12	0	0	0	0	0	12	25	100	12.7	2	0.2	4	0.4	6	0.6
		後	17	0	0	0	0	0	17	17	100	13.9	2	0.2	5	0.6	7	0.8

調査時期			セジロウシ									
年	月	期	調査発生		寄生株率		株当成虫数		株当幼虫数		合計虫数	
			調査数	発生数	最高	平均	最高	平均	最高	平均	最高	平均
98	5	前	20	10	4	0.4	-	-	-	-	-	-
		後	42	23.3	20	1.9	0.8	0.04	0.2	0.005	0.8	0.05
	6	前	15	80	44	14.9	4.2	0.3	4.2	0.4	8.4	0.8
		後	38	78.9	100	37.4	3	0.5	20	1.8	22.5	2.3
	7	前	13	69.2	100	50.2	1	0.1	20	4.6	20.2	4.7
		後	33	54.5	100	27.6	2.6	0.3	5	0.6	6.3	0.9
	8	前	12	75	100	35.5	10	1.5	6.2	0.8	11.6	2.3
		後	17	5.9	100	5.9	5	0.3	3	0.2	8	0.5

D カンキツのチャノキイロアザミウマ
(黄色粘着トラップ)

月	半 旬	チャバネアオカメムシ						ツヤアオカメムシ					
		疎 早		島原	佐世保	岐宿	西彼	疎 早		島原	佐世保	岐宿	西彼
		白熱	青色	白熱	白熱	白熱	青色	白熱	青色	白熱	白熱	白熱	青色
4	1			-	-	-	-			-	-	-	-
	2			-	-	-	-			-	-	-	-
	3			-	-	-	-		1	-	-	-	-
	4		2	-	-	-	-		1	-	-	-	-
	5		25	-	-	-	-		39	-	-	-	-
	6	7	12	-	-	-	-	1	8	-	-	-	-
5	1		74	-	-	-		2	104	-	-	-	
	2		98	-	-	-			315	-	-	-	
	3		1	-	-	-		1	11	-	-	-	
	4		33	-	-	-	29		66	-	-	-	14
	5		17	-	-	-	30		29	-	-	-	2
	6	1	16	-	-	-	21	1	18	-	-	-	7
6	1		7		1	1	7		13		1	2	3
	2		7	1			1		5			1	2
	3		22		1	10	51		17			27	11
	4		34	1			41		13			3	8
	5		32		* 2		135		8			2	20
	6	1	27	1		10			15	1	2	29	
7	1	1	26			7			1		1	19	
	2		26				172						33
	3		36				194				2		16
	4		39		1		63					1	26
	5		32		1		243	1	4		1		50
	6	1	25	1	5	1	172		7		2	1	27
8	1		76	3	7		353		22	3	19		48
	2	2	196	1	9		55		41		3		8
	3	4	85	8	25			2	35	3	21	1	
	4	4	43	13	30		29		12	8	64	1	9
	5		12	5	15		40		5	1	29		2
	6		43	3	23		11		77	4	20		
9	1		14	1	10			1	28	2	14		
	2		20	1	83				57		12		
	3	1	16		61		73	1	25	2	107		4
	4		9	2	115		25	2	30	2	159		4
	5		35	4	25		10		17	6	72		1
	6		17		17		90		12		27		10
10	1		27	-	-	-	3		23	-	-	-	2
	2		7	-	-	-	2		9	-	-	-	2
	3	1	7	-	-	-	9	1	20	-	-	-	11
	4			-	-	-	1		1	-	-	-	
	5		1	-	-	-			4	-	-	-	1
	6			-	-	-			1	-	-	-	1

調査時期	県予察圃場
98. 4. 1	設 置
10	0
18	0
27	1
5. 1	5
8	5
15	3
29	7
6. 5	29
12	41
19	11
26	45
7. 3	19
10	13
17	41
24	41
31	36
8. 7	67
14	84
21	107
28	137
9. 4	145
11	183
18	162
25	97
10. 2	177
9	59
16	17
23	4
30	0
合 計	1,537

※ 表中の-は未調査

佐世保(白熱)の6月5半旬のデータは4～5半旬の合計値

Ｃ カメムシ類

月	半 旬	クモヘリカメムシ					ホソハリカメムシ					シラホシカメムシ					アカスジメクラガメ				
		諫 早		島原	佐世保	岐宿	諫 早		島原	佐世保	岐宿	諫 早		島原	佐世保	岐宿	諫 早		島原	佐世保	岐宿
		白熱	青色	白熱	白熱	白熱	白熱	青色	白熱	白熱	白熱	白熱	青色	白熱	白熱	白熱	白熱	青色	白熱	白熱	白熱
4	1			-	-	-			-	-	-			-	-	-			-	-	-
	2			-	-	-			-	-	-			-	-	-			-	-	-
	3			-	-	-			-	-	-			-	-	-			-	-	-
	4			-	-	-			-	-	-			-	-	-			-	-	-
	5			-	-	-			-	-	-			-	-	-			-	-	-
	6			-	-	-			-	-	-			-	-	-			-	-	-
5	1			-	-	-			-	-	-			-	-	-	2		-	-	-
	2			-	-	-			-	-	-			-	-	-			-	-	-
	3			-	-	-			-	-	-			-	-	-	5		-	-	-
	4			-	-	-			-	-	-			-	-	-			-	-	-
	5			-	-	-			-	-	-			-	-	-	6		-	-	-
	6			-	-	-		1	-	-	-			-	-	-	1		-	-	-
6	1						1										16	2			
	2												2				1				
	3							1									21				
	4																7				
	5	1															2				
	6	1	1														11				
7	1	2						1													
	2	3																			
	3	7																			
	4	15				1										1					
	5	7	1																		
	6	6	1			5															
8	1		3			3		15													
	2	1		1	1			1								1					
	3		1	1	1			12					3			2		1			
	4		1	1	1			5					1					1			
	5		3		1			1	1							1					
	6	4	14	7				1				1									
9	1		8	5	1			1													
	2	1	1	5	3			2													
	3		1	2	1			3													
	4			1	2													1			
	5		1		3																
	6							1													
10	1		1	-	-	-			-	-	-			-	-	-			-	-	-
	2			-	-	-			-	-	-			-	-	-			-	-	-
	3			-	-	-			-	-	-			-	-	-			-	-	-
	4		1	-	-	-			-	-	-			-	-	-			-	-	-
	5			-	-	-			-	-	-			-	-	-			-	-	-
	6			-	-	-			-	-	-			-	-	-			-	-	-

※ 表中の-は未調査

B. ハスモンヨトウ、ナシヒメシシクイ、チャノコカクモンハマキ、チャノホソガ

月	日	ハスモンヨトウ		ナシヒメシシクイ		チャノコカクモンハマキ				チャノホソガ
		諫早(フェロモントラップ)		諫早(フェロモントラップ)		東彼杵(予察灯、白熱)		東彼杵(フェロモントラップ)		東彼杵(フェロモントラップ)
		1998	平 年	1998	平 年	1998	平 年	1998	平 年	1998
4	1	1	1.1	9	5.0	0	0	0	3.0	0
	2	2	3.1	5	7.0	0	0.2	11	3.8	20
	3	8	2.6	5	2.8	0	0	5	17.2	22
	4	11	4.9	2	5.3	0	0.2	7	18.0	13
	5	7	11.8	2	5.0	1	0.2	31	93.0	11
	6	10	26.9	1	1.3	0	6.8	35	229.0	0
5	1	56	38.3	0	0.7	0	4.6	20	98.0	2
	2	47	29.3	1	1.0	0	13.4	0	354.0	0
	3	72	30.6	4	1.7	0	3.2	4	227.0	1
	4	65	41.7	6	1.4	0	9.4	0	327.0	39
	5	27	36	8	0.7	0	1.8	1	222.0	484
	6	30	56.9	11	5.3	0	2.4	1	39.2	404
6	1	43	73.7	2	9.0	1	0	29	2.8	408
	2	12	76	4	6.6	0	0.4	122	4.4	112
	3	31	93.9	4	4.3	1	0.2	110	13.0	228
	4	128	138.7	12	5.9	0	2.6	57	170.0	123
	5	142	144	9	5.1	4	32.8	146	262.0	56
	6	91	183.9	5	6.4	1	69.8	31	177.0	0
7	1	49	196.2	30	9.4	2	44.0	8	222.0	73
	2	9	144.6	11	21.0	0	42.8	9	183.0	361
	3	15	155.6	16	17.7	0	4.2	108	77.8	262
	4	50	160.1	11	17.1	3	0.6	37	19.2	56
	5	102	115.4	16	9.9	10	1.6	132	318.0	197
	6	108	149.1	20	13.0	2	1.4	63	233.0	490
8	1	77	162.9	15	10.3	0	17.6	47	354.0	602
	2	59	173	14	11.1	0	110.0	56	567.0	79
	3	65	183.9	12	11.6	0	122.0	16	798.0	116
	4	218	195.9	22	13.9	0	27.0	108	133.0	107
	5	189	236.9	18	5.1	0	3.0	151	39.6	102
	6	159	332.9	19	5.1	3	1.6	341	25.8	764
9	1	164	354.4	25	7.8	1	0.8	96	96.0	966
	2	72	365	30	11.2	0	2.2	31	202.0	473
	3	94	429.6	20	8.8	0	5.4	36	396.0	962
	4	201	444.7	5	7.6	0	8.6	7	575.0	223
	5	106	451.1	1	4.4	0	6.6	27	547.0	706
	6	228	464.4	9	2.6	0	7.2	10	260.0	190
10	1	178	392.9	5	4.5	0	0.4	65	71.2	1,180
	2	189	517.9	0	2.3	0	0.4	26	37.4	1,189
	3	154	486.4	1	0.5	0	1.0	80	27.2	702
	4	694	498.8	0	0.0	0	0.4	187	23.8	1,403
	5	300	356.1	1	0.0	0	0.6	51	9.4	422
	6	205	441.8	0	0.0	0	0	124	7.8	1,072
11	1	184	289	-	-	-	-	-	-	-
	2	86	212	-	-	-	-	-	-	-
	3	125	175.5	-	-	-	-	-	-	-
	4	60	121	-	-	-	-	-	-	-
	5	7	98	-	-	-	-	-	-	-
	6	19	126	-	-	-	-	-	-	-
合 計		4,949	9,424.2	391	270	29	557.4	2,426	7,483.6	14,620

※ チャノコカクモンハマキ、チャノホソガは総合農林試験場東彼杵茶葉支場の調査