

2 発生面積及び防除面積

(単位: ha, %)

農作物名	作付面積	病害虫名	程度別発生面積					防除面積			
			甚	多	中	少	計	実	延		
水稲 (早期)	2,000	苗いもち				3	3	*	1,700	*	1,700
		葉いもち		2	13	715	730		1,300		1,400
		穂いもち			5	385	390		1,800		1,800
		紋枯病			20	620	640		1,800		1,800
		白葉枯病					0		0		0
		ばか苗病				50	50		1,300		1,300
		もみ枯細菌病				10	10		0		0
		ごま葉枯病			30	610	640	*	1,700	*	1,700
		黄化萎縮病					0		0		0
		縞葉枯病				200	200		-		-
		萎縮病					0		-		-
		ニカメイガ					0		0		0
		セジロウンカ				2,000	2,000	*	1,800	*	2,700
		トビイロウンカ		10	20	1,410	1,440	*	1,800	*	2,700
		ヒメトビウンカ				1,760	1,760	*	1,800	*	2,700
		ツマグロヨコバイ				1,680	1,680	*	1,800	*	2,700
		イネゾウムシ			3	477	480	*	1,700	*	1,700
		イネクロカメムシ			20	270	290		100		100
		斑点米カメムシ類			90	670	760		1,800		3,300
		イチモンジセセリ				160	160	*	1,800	*	1,800
		フタオビコヤガ				240	240	*	1,800	*	1,800
		イネヨトウ				20	20	*	1,800	*	1,800
		アワヨトウ				20	20	*	1,800	*	1,800
		コブノメイガ			40	1,520	1,560		1,800		1,800
		イネミズゾウムシ			45	1,235	1,280		1,700		1,800
水稲 (普通期)	11,600	苗いもち			1	29	30	*	7,500	*	7,500
		葉いもち		70	200	2,070	2,340		10,400		10,400
		穂いもち		10	50	2,140	2,200		10,000		10,000
		紋枯病		10	50	3,935	3,995		10,200		10,200
		白葉枯病					0				
		ばか苗病			5	123	128		7,400		7,400
		もみ枯細菌病		10	50	2,040	2,100		1,500		1,500
		ごま葉枯病		30	100	2,220	2,350	*	7,500	*	7,500
		黄化萎縮病					0		0		0
		縞葉枯病				1,610	1,610		-		-
		萎縮病					0		-		-
		ニカメイガ					0		0		0
		セジロウンカ				8,470	8,470	*	11,000	*	30,000
		トビイロウンカ	100	200	350	10,550	11,200	*	11,000	*	30,000
		ヒメトビウンカ				9,400	9,400	*	11,000	*	30,000
		ツマグロヨコバイ				4,840	4,840	*	11,000	*	30,000
		イネゾウムシ			2	288	290	*	8,000	*	8,000
		イネクロカメムシ		1	10	509	520		250		250
		斑点米カメムシ類		5	250	3,935	4,190		10,700		16,000
		イチモンジセセリ				460	460	*	10,500	*	19,000
		フタオビコヤガ			2	818	820	*	10,500	*	19,000
		イネヨトウ				50	50	*	10,500	*	19,000
		アワヨトウ				30	30	*	10,500	*	19,000
		コブノメイガ			50	5,100	5,150		10,500		18,000
		イネミズゾウムシ			30	1,130	1,160		8,000		8,100
スクミリンゴガイ	20	600	600	2,580	3,800		1,600		1,600		
麦	1,747	さび病類				140	140		0		0
		うどんこ病				118	118	*	1,700	*	3,300
		赤かび病				165	165		1,700		3,300
		黒穂病類			1	364	365		300		300
		斑葉病				16	16	*	300	*	300
		小麦縞萎縮病					0		0		0
		大麦縞萎縮病					0		0		0
		株腐病				30	30		0		0
		アブラムシ類			10	1,290	1,300		25		25
		アブラムシ類									
大豆	404	紫斑病			5	45	50		100		150
		べと病			5	355	360		0		0
		葉焼病			10	350	360		0		0
		モザイク病					0				
		アブラムシ類				130	130	*	300	*	300
		コガネムシ類				20	20	*	30	*	30
		ハスモンヨトウ			10	290	300		300		1,000
		ダイズサヤタマバエ				50	50	*	380	*	1,200
		ヒメサヤムシ類				30	30	*	380	*	1,200
		マメシンクイガ				40	40	*	380	*	1,200
		シロイチモジマダラメイガ				40	40	*	380	*	1,200
		ウワバ類				90	90	*	300	*	1,000
		カメムシ類			30	170	200	*	380	*	1,000
ばれいしょ	4,220 春作:3,040 秋作:1,180	疫病			10	90	100		3,050		7,600
		葉巻病				0	0		-		-
		アブラムシ類			75	2,925	3,000		3,000		5,100
		ニジュウヤホシテントウ				190	190	*	3,000	*	5,100
		食葉性ヤガ類			47	520	567	*	3,000	*	5,100

(単位: ha, %)									
農作物名	作付面積	病害虫名	程度別発生面積					防除面積	
			甚	多	中	少	計	実	延
冬春きゅうり (2012年植付)	42	モザイク病				1	1	-	-
		斑点細菌病	1	1	2	1	1	10	15
		べと病				34	38	42	210
		褐斑病				9	10	42	170
		うどんこ病				30	37	42	165
		炭疽病				1	1	5	10
		灰色かび病				5	5	42	140
		菌核病				3	3	42	140
冬春トマト (2012年植付)	117	アブラムシ類				5	5	* 42	* 60
		ミミキロアザミウマ			4	20	24	42	160
		疫病			1	10	11	117	205
		灰色かび病			8	31	39	117	610
		モザイク病				1	1	-	-
たまねぎ (2012年植付)	677	コナジラミ類			1	66	67	117	700
		アブラムシ類				10	10	* 117	* 130
		白色疫病	1	1	13	30	30	677	1,700
		べと病				210	225	677	1,800
		ホトリキス葉枯病				195	230	-	-
冬レタス (2012年植付)	590	ネギアザミウマ			20	585	605	677	1,350
		灰色かび病		3	10	30	40	295	550
		菌核病			27	190	220	* 295	* 550
		斑点細菌病			4	36	40	295	550
		腐敗病				20	20	* 295	* 550
いちご (2012年植付)	284	アブラムシ類			3	82	85	590	* 1,350
		ネキリムシ類				15	15	270	270
		ハスモンヨトウ			10	130	140	590	1,350
		オオタバコガ				40	40	* 590	* 1,350
		炭疽病			2	32	34	284	2,300
かんきつ	3,933	萎黄病			1	14	15	230	230
		灰色かび病		1	8	74	83	284	940
		うどんこ病			9	21	30	284	2,200
		アブラムシ類	1	1	4	95	100	284	490
		ハダニ類		2	20	133	156	284	1,130
びわ	584	ハスモンヨトウ		1	9	150	160	284	850
		そうか病			20	200	220	3,800	7,600
		かいよう病			60	400	460	1,500	2,000
		黒点病		30	320	2,800	3,150	3,900	15,600
		灰色かび病			90	890	980	* 3,900	* 4,700
		ミカンハダニ	1	15	240	2,350	2,606	3,900	7,800
		ミカンサビダニ		1	24	255	280	* 3,900	* 7,800
		チャノキイロアザミウマ		5	200	1,000	1,205	3,800	11,400
		カメムシ類		1	48	400	450	1,500	2,500
		ヤノネカイガラムシ	1	1	5	100	105	* 2,900	* 4,000
		マルカイガラムシ類			25	250	275	* 2,600	* 4,000
		ロウムシ類			2	38	40	* 2,600	* 4,000
		ゴマダラカミキリ		5	90	150	245	* 1,500	* 2,500
		クワゴマダラヒトリ			5	200	205	200	200
		アブラムシ類			100	1,430	1,530	* 3,900	* 7,800
なし	73	がんしゅ病	0	5	30	435	470	410	630
		灰斑病	0	25	120	439	584	450	670
		ナシヒメシンクイ	0	2	10	350	362	350	400
		カメムシ類	0	0	0	20	20	2	2
ぶどう	101	黒星病		2	13	25	40	73	550
		赤星病			2	38	40	73	150
		うどんこ病			1	29	30	* 73	* 120
		輪紋病			1	14	15	73	150
		ナシヒメシンクイ			5	35	40	73	150
		ハダニ類			12	38	50	60	100
		カメムシ類			10	10	10	25	75
		アブラムシ類		2	5	33	40	73	200
茶	753	黒とう病			2	33	35	101	450
		べと病			3	32	35	101	400
		うどんこ病				5	5	* 100	* 200
		晩腐病			3	10	13	101	300
		褐斑病				10	10	* 101	* 210
		灰色かび病			2	16	18	101	210
		チャノキイロアザミウマ			10	30	40	101	420
		炭そ病			30	470	500	750	1,500
茶	753	もち病				20	20	* 360	* 720
		チャノコカクモンハマキ		10	60	380	450	750	2,250
		チャノホソガ		10	60	410	480	* 750	* 2,250
		チャノミドリヒメヨコバイ		20	80	550	650	* 750	* 2,250
		カンザワハダニ		5	30	400	435	500	1,500
		チャノキイロアザミウマ	2	18	110	540	670	750	2,250
		クワシロカイガラムシ		10	90	440	540	750	1,400

注) 防除面積の*は同時防除